

## SOME NEW HALTICINI FROM INDIANA AND ONTARIO

BY W. S. BLATCHLEY,

INDIANAPOLIS, INDIANA.

Since my paper entitled "Notes on Indiana Halticini, etc." was published in this JOURNAL<sup>1</sup>, a number of additional species have been taken in Southern Indiana, four of which are evidently new to science. Among some Coleoptera sent to me for naming by J. F. Brimley, of Wellington, Ontario, was also an undescribed species of *Longitarsus*. The descriptions of these new forms are as follows:

### *Disonycha admirabila*, new species.

Oblong, subparallel, convex. Thorax, under surface and femora dull yellow; antennæ in great part, scutellum and a sutural discal and submarginal stripe on each elytron, black; head and two narrow stripes on each elytron dull red, the red stripes bordered each side by a narrow one of silvery-white which lies adjacent to the black stripe; tibiæ and tarsi wholly fuscous-black; labrum and basal joint of antennæ piceous; last joint of antennæ pale. Head impunctate, the tubercles and frontal carina small, obtuse. Thorax two-thirds wider than long, absolutely smooth, sides rounded into base; front margin truncate, hind one convex. Elytra slightly wider at base than thorax, very finely alutaceous, minutely and very sparsely punctate. Abdomen and femora very finely punctate, minutely pubescent. Length 4.5 mm.

Knox Co., Ind., September 18. Taken by sweeping smartweed (*Polygonum*) in a dry upland woods. Allied to *caroliniana* (Fab.) but differs in its smaller more parallel form, dark labrum, wholly dark tibiæ and the alutaceous tricolored elytra. From *crenicollis* (Say) it differs in its smaller size, much narrower non-alutaceous and impunctate thorax without black spots and much more finely alutaceous elytra.

### *Longitarsus pallescens*, new species.

Broadly oval, convex, robust for the genus. Color above a uniform pale dull yellow; apical half of antennæ brown, labrum shining black; abdomen in great part chestnut-brown. Head impunctate, occiput finely alutaceous; antennæ very slender, joints 2-4 gradually slightly increasing in length. Thorax subquadrate, one-third wider than long, sides rather broadly rounded, front angles thickened, disk rugose, rather coarsely, shal-

<sup>1</sup>Vol. XIX, No. 1, March, 1921, pp. 16-27.

lowly irregularly punctate. Elytra oval, one-third wider at base than thorax, humeri rounded, umbones evident but small; disk finely, very shallowly evenly punctate. Under surface and femora very finely punctate, minutely pubescent. Length 2.8 mm.

Prince Edward County, Ontario, April 7. Two specimens collected by J. F. Brimley. Belongs under *c*, p. 18 of my key, *loc. cit.*, and allied to *subrufus* Lec., but differs in its pale color, much broader form, more slender antennæ, alutaceous occiput and rugosely punctate thorax.

***Longitarsus acutipennis*, new species.**

Elongate, convex; widest at middle of abdomen, strongly tapering behind, color a uniform shining pale brownish-yellow. Head impunctate, occiput minutely transversely strigose. Antennæ slender, three-fourths the length of body, joints 2-4 subequal in length. Thorax subelliptical, convex, one-half wider than long, sides rounded; disk minutely alutaceous, very finely and sparsely unevenly punctate. Elytra narrower at base than middle of thorax, thence gradually widened to middle, then tapering to the narrowly rounded or subacute tips; humeri, umbones and inner wings absent; pygidium concealed; disk rather coarsely, closely and evenly punctate. Length 1.8 mm.

Crawford County, Ind., August 5. Swept from goldenrod. Belongs under *pp*, p. 19 of my key, except that the pygidium is concealed. Of the twenty-one eastern species of *Longitarsus* at hand this differs from all in the strongly tapering posterior portion of the body. It is also paler than any of those belonging to the subgenus *Apterius* in which the humeri, umbones and inner wings are wanting.

***Glyptina ferruginea*, new species.**

Broadly oval, robust. Dark reddish-brown, shining, thorax and under surface somewhat paler than elytra; apical half of antennæ fuscous-brown. Head smooth. Antennæ slender, half the length of body, joints 2-4 subequal in length, 2 stouter than 3. Thorax subquadrate, convex, less than one-half wider than long, not narrowed in front, sides feebly curved, front angles slightly obliquely truncate; disk very finely, sparsely aciculate punctate. Elytra at base nearly one-half wider than thorax, conjointly oval, tips broadly rounded, striæ coarsely, somewhat rugosely and closely punctate, the ninth stria not more impressed than those on disk; intervals almost flat, each with a single row of very minute punctures. Length 1.3-1.5 mm.

Crawford Co., Indiana, August 1-4; three specimens taken by sweeping low herbage along a pathway on high wooded slopes. Allied to *brunnea* Horn but distinctly shorter and relatively broader, with antennæ darker and punctures of elytra coarser.

**Glyptina leptosoma**, new species.

Narrowly oval, strongly convex, tapering behind. Above and beneath reddish-brown, shining; antennæ reddish-brown, the apical third fuscous; legs pale brownish-yellow. Antennæ slender, two-thirds the length of body, second joint not stouter than third, slightly more than two-thirds the length of latter. Head impunctate. Thorax subquadrate, strongly convex, scarcely wider than long, side margins broadly curved, their front ends tuberculate; disk rather coarsely, shallowly, not densely punctate. Elytra conjointly oval, about one-fourth wider at base than thorax, sides broadly curved, strongly tapering behind the middle, to the narrowly rounded tips; disk strongly convex, punctures of striæ relatively coarse, those on sides finer; intervals flat, impunctate. Abdomen finely and sparsely punctate, minutely pubescent. Length 1.2 mm.

Crawford County, Ind., August 4; taken by sweeping goldenrod. Smaller, narrower and more strongly convex than either *brunnea* or *feruginea*, the thoracic punctures coarser and the elytra much more narrowed behind the middle. In form it much resembles some of the smaller species of *Longitarsus*.